

Annex 1: List of most important publications of the NCCR

- Ayyanan, A., Civenni, G., Ciarloni, L., Morel, C., Lefort, K., Mandinova, A., Raffoul, W., Fiche, M., Dotto, G. P., and Brisken, C. (2006). Increased Wnt signaling triggers oncogenic conversion of human mammary epithelial cells by a Notch-dependent mechanism in mammary gland development. *Proc. Natl. Acad. Sci. USA* **103**, 3799-3804.
- Bossen, C., Cachero, T.G., Tardivel, A., Ingold, K., Willen, L., Dobles, M., Scott, M.L., Maquetin, A., [...], Tschopp, J., and Schneider, P. (2008). TACI, unlike BAFF-R, is solely activated by oligomeric BAFF and APRIL to support survival of activated B cells and plasmablasts. *Blood* **111**, 1004-1012.
- Ciarloni, L., Mallepell, S., and Brisken, C. (2007). Amphiregulin is an essential mediator of estrogen receptor alpha function in mammary gland development. *Proc. Natl. Acad. Sci. USA* **104**, 5455-5460.
- Corbett, M., Boger, W., Heeney, J.L., Gerber, S., Genin, C., Didierlaurent, A., Oostermeijer, H., Dubbes, R., Braskamp, G., Lerondel, S., et al. (2008). Aerosol immunization with NYVAC and MVA vectored vaccines is safe, simple, and immunogenic. *Proc. Natl. Acad. Sci. USA* **105**, 2046-2051.
- De Vito C, Riggi N, Cornaz S, Suva ML, Baumer K, Provero P, Stamenkovic I. (2012). A TARBP2-dependent miRNA expression profile determines cancer stem cell properties and provides candidate therapeutic reagents in Ewing sarcoma. *Cancer Cell* **21**, 807-821.
- Dostert, C., Petrilli, V., Van Bruggen, R., Steele, C., Mossman, B.T., and Tschopp, J. (2008). Innate immune activation through Nalp3 inflammasome sensing of asbestos and silica. *Science* **320**, 674-677.
- Farmer, P., Bonnefoi, H., Anderle, P., Cameron, D., Wirapati, P., Becette, V., André, S., Piccart, M., Campone, M., Brain, E., MacGrogan, G., Petit T, Jassem J, Bibeau, F., Blot, E., Bogaerts, J., Aguet, M., Bergh, J., Iggo, R., and Delorenzi, M. (2009). A stroma-related gene signature predicts resistance to epirubicin-containing neoadjuvant chemotherapy in breast cancer. *Nat. Med.* **15**, 68-74.
- Farmer, P., Bonnefoi, H., Becette, V., Tubiana-Hulin, M., Fumoleau, P., Larsimont, D., MacGrogan, G., Bergh, J., Cameron, D., Goldstein, D., Duss, S., Nicoulaz, A.L., Brisken, C., Fiche, M., Delorenzi, M., and Iggo, R. (2005). Identification of molecular apocrine breast tumours by microarray analysis. *Oncogene* **24**, 4660-4671.
- Fevr, T., Robine, S., Louvard, D., and Huelsken, J. (2007). Wnt/b-catenin is essential for intestinal homeostasis and maintenance of intestinal stem cells. *Mol. Cell Biol.* **27**, 7551-7559.
- Hegi, M.E., Diserens, A.C., Gorlia, T., Hamou, M.F., de Tribolet, N., Weller, M., Kros, J.M., Hainfellner, J.A., Mason, W., Mariani, L., et al. (2005). MGMT gene silencing and benefit from temozolomide in glioblastoma. *N. Engl. J. Med.* **352**, 997-1003.
- Ingold, K., Zumsteg, A., Tardivel, A., Huard, B., Steiner, Q.G., Cachero, T.G., Qiang, F., [...], Tschopp, J., and Schneider, P. (2005). Identification of proteoglycans as the APRIL-specific binding partners. *J. Exp. Med.* **201**, 1375-1383.
- Jeannet, G., Scheller, M., Scarpellino, L., Duboux, S., Gardiol, N., Back, J., Kuttler, F., Malanchi, I., Birchmeier, W., Leutz, A., Huelsken*, J., and Held*, W. (2008). Long-term, multilineage hematopoiesis occurs in the combined absence of b-catenin and g-catenin. *Blood* **111**, 142-9.
*these two authors contributed equally.
- Malanchi, I., Peinado, H., Kassen, D., Husseinet, T., Metzger, D., Champon, P., Huber, M., Hohl, D., Cano, A., Birchmeier, W., and Huelsken, J. (2008). Cutaneous cancer stem cell maintenance is dependent on b-catenin signaling. *Nature* **452**, 650-3.

Malanchi*, I., Santamaria-Martínez*, A., Susanto, E., Peng, H., Lehr, H.A., Delaloye, J.F., and Huelsken, J. (2012). Interactions between cancer stem cells and their niche govern metastatic colonization. *Nature* 481, 85–89. *these two authors contributed equally

Murat, A., Migliavacca, E., Gorlia, T., Lambiv, W.L., Shay, T., Hamou, M.-F., de Tribolet, N., Regli, L., Wick, W., Kouwenhoven, M.C.M., Hainfellner, J.A., Heppner, F.L., Dietrich, P.-Y., Zimmer, Y., Cairncross, J. G., Janzer, R.-C., Domany, E., Delorenzi, M., Stupp, R., and Hegi, M.E. (2008). Stem Cell-Related "Self-Renewal Signature" and High Epidermal Growth Factor Receptor Expression Associated with Resistance to Concomitant Chemoradiotherapy in Glioblastoma. *J. Clin. Oncol.* 26, 3015-3024.

Naef, F., and Huelsken, J. (2005). Cell-type-specific transcriptomics in chimeric models using transcriptome-based masks. *Nucleic Acids Res.* 33, e111.

Riggi, N., Cironi, L., Provero P., Suvà, M.L., Kaloulis, K., Garcia-Echeverria, C., Hoffmann, F., Trumpp, A. and Stamenkovic, I. (2005). Development of Ewing's sarcoma from bone-marrow-derived mesenchymal progenitor cells. *Cancer Res.* 65, 11459-1468.

Riggi, N., Suvà, M.L., De Vito, C., Provero, P., Stehle, J.C., Baumer, K., Cironi, L., Janiszewska, M., Petricevic, T., Suvà, D., Tercier, S., Joseph, J.M., Guillou, L., and Stamenkovic, I. (2010). EWS-FLI-1 modulates miRNA145 and SOX2 expression to initiate mesenchymal stem cell reprogramming toward Ewing sarcoma cancer stem cells. *Genes Dev.* 24, 916-932.

Rufer, N., Zippelius, A., Batard, P., Pittet, M.J., Kurth, I., Corthesy, P., Cerottini, J.C., Leyvraz, S., Roosnek, E., Nabholz, M., et al. (2003). Ex vivo characterization of human CD8+ T subsets with distinct replicative history and partial effector functions. *Blood* 102, 1779-1787.

Sabine, A., Agalarov, Y., Maby-EI Hajjami, H., Jaquet M., Hägerling, R., Pollmann, C., Bebber, D., Pfenniger, A., Miura, N., Dormond, O., Calmes, J.-M., Adams, R.H., Mäkinen, T., Kiefer, B., Kwak, B.R., and Petrova, T.V (2012). Mechanotransduction, PROX1 and FOXC2 cooperate to control connexin37 and calcineurin during lymphatic valve formation. *Dev. Cell* 22, 420-445

Schmid, D.A., Irving, M.B., Posevitz, V., Hebeisen, M., Posevitz-Fejfar, A., Sarria, J.-C.F., Gomez-Eerland, R., Thome, M., Schumacher, T.N.M., Romero, P., et al. (2010). Evidence for a TCR affinity threshold delimiting maximal CD8 T cell function. *J. Immunol.* 184, 4936–4946.

Sotiriou C, Wirapati P, Loi S, Harris A, Bergh J, Smeds J, Farmer P, Praz V, Haibe-Kains B, Desmedt C, Larsimont D, Cardoso F, Peterse H, Nuyten D, Buyse M, Van de Vijver MJ, Piccart M, Delorenzi M. (2006). Gene expression profiling in breast cancer: understanding the molecular basis of histologic grade to improve prognosis. *J. Natl. Cancer Inst.* 98, 262-272.

Wicki, A., Lehembre, F., Wick, N., Hantusch, B., Kerjaschki D., and Christofori, G. (2006). Tumor invasion in the absence of epithelial-mesenchymal transition: podoplanin-mediated remodeling of the cytoskeleton. *Cancer Cell* 9, 261-272.

Zippelius, A., Pittet, M.J., Batard, P., Rufer, N., de Smedt, M., Guillaume, P., Ellefsen, K., Valmori, D., Lienard, D., Plum, J., et al. (2002). Thymic selection generates a large T cell pool recognizing a self-peptide in humans. *J. Exp. Med.* 195, 485-494.

Zaric, J., Tercier, S., Joseph, J.M., Sengstag, T., Delorenzi, M., Ponsonnet, L., and Rüegg, C. (2012). Identification of MAGI1 as a tumor suppressor protein induced by COX-2 inhibitors in colon cancer cells. *Oncogene* 31, 48–59.